


Best Practice in Perineal Care

[View Online](#)

Aasheim, V., Nilsen, A. B. V., Lukasse, M., & Reinar, L. M. (1996). Perineal techniques during the second stage of labour for reducing perineal trauma. Cochrane Database of Systematic Reviews. <http://doi.wiley.com/10.1002/14651858.CD006672.pub2>

Albers, L...[et al]. (2005). Midwifery care measures in the second stage of labor and reduction of genital tract trauma at birth: a randomised trial. *Journal of Midwifery & Women's Health*, 50(5).

Albers, L...[et al]. (2006). Body mass index, midwifery intrapartum care, and childbirth lacerations. *Journal of Midwifery & Women's Health*, 51(4).

Albers, L...[et al]. (2006). Factors related to genital tract trauma in normal spontaneous vaginal births. *Birth*, 33(2).

Andrews, Gilly. (2005). *Women's sexual health* (3rd ed). Elsevier.

Andrews, V...[et al]. (2006). Occult anal sphincter injuries - myth or reality? *British Journal of Obstetrics and Gynaecology*, 113(2).

Beckmann, M. M. and A. J. Garratt. (2006). Antenatal perineal massage for reducing perineal trauma. *The Cochrane Database of Systematic Reviews*, Issue 1.

Bick, D. E., Ismail, K. M., Macdonald, S., Thomas, P., Tohill, S., & Kettle, C. (2012). How good are we at implementing evidence to support the management of birth related perineal trauma? A UK wide survey of midwifery practice. *BMC Pregnancy and Childbirth*, 12(1). <https://doi.org/10.1186/1471-2393-12-57>

Bick, D. E., Kettle, C., Macdonald, S., Thomas, P. W., Hills, R. K., & Ismail, K. M. (2010). PERineal Assessment and Repair Longitudinal Study (PEARLS): protocol for a matched pair cluster trial. *BMC Pregnancy and Childbirth*, 10(1). <https://doi.org/10.1186/1471-2393-10-10>

Bick, Debra, MacArthur, Christine, Winter, Heather, Mott, Kathy de, & Ritchie, Gill. (2009). *Postnatal care: evidence and guidelines for management* (2nd ed). Churchill Livingstone Elsevier.

Borders, N. (2006). After the afterbirth: A critical review of postpartum health relative to method of delivery. *Journal of Midwifery & Women's Health*, 51(4).

Boyle, Maureen. (2006). *Wound healing in midwifery*. Radcliffe Publishing.

- Caroci, A. S. and M. L. G. Riesco. (2006). A comparison of 'hands off' versus 'hands on' techniques for decreasing perineal lacerations during birth. *Journal of Midwifery & Women's Health*, 51(2).
- Carroli, G. and L. Mignini. (2009). Episiotomy for vaginal birth. *The Cochrane Database of Systematic Reviews*, Issue 1.
- Chapman, V. (2009). Perineal trauma and suturing. In *The midwife's labour and birth handbook* (2nd ed). Wiley-Blackwell.
- Chapman, V. (2009). Clinical skills: issues affecting the left-handed midwife. *British Journal of Midwifery*, 17(9), 588–592. <https://doi.org/10.12968/bjom.2009.17.9.43900>
- Dudding, T. C., Vaizey, C. J., & Camm, M. A. (2008). Obstetric anal sphincter injury: incidence, risk factors and management (Review). *Annals of Surgery*, 247(2), 224–237.
- East, C. (2012). Perineal pain following childbirth: prevalence,. *Midwifery*, 28(1), 93–97. <http://www.ncbi.nlm.nih.gov/pubmed/21236531>
- Eliminating female genital mutilation: an interagency statement. (n.d.).
- Female genital mutilation. (n.d.).
- Fernando, R. J. (2007). Risk factors and management of obstetric perineal injury. *Obstetrics, Gynaecology & Reproductive Medicine*, 17(8).
- Fleming, N., E. R. Newton and J. Roberts. (2003). Changes in postpartum perineal muscle function in women with and without episiotomies. *Journal of Midwifery & Women's Health*, 48(1).
- Fornell, E. U. (2005). Obstetric anal sphincter injury ten years after: subjective and objective long term effects. *BJOG: An International Journal of Obstetrics and Gynaecology*, 112(3), 312–316. <https://doi.org/10.1111/j.1471-0528.2004.00400.x>
- Groom, K. M. and S. Paterson-Brown. (2002). Can we improve on the diagnosis of third degree tears? *European Journal of Obstetrics, Gynecology and Reproductive Biology*, 101(1).
- Gupta, J. K., G. J. Hofmeyr and R. Smyth. (2004). Position in the second stage of labour for women without epidural anaesthesia. *The Cochrane Database of Systematic Reviews*, Issue 1.
- Gurol-Urganci, I., Cromwell, D., Edozien, L., Mahmood, T., Adams, E., Richmond, D., Templeton, A., & van der Meulen, J. (2013). Third- and fourth-degree perineal tears among primiparous women in England between 2000 and 2012: time trends and risk factors. *BJOG: An International Journal of Obstetrics & Gynaecology*. <https://doi.org/10.1111/1471-0528.12363>
- Hartmann, K...[et al]. (2005). Outcomes of routine episiotomy: A systematic review. *JAMA the Journal of the American Medical Association*, 293(17).
- Hedayati, H., J. Parsons, and C. A. Crowther. (2003). Rectal analgesia for pain from perineal

trauma following childbirth. The Cochrane Database of Systematic Reviews, Issue 3.

Henderson, Christine & Bick, Debra. (2005). Perineal care: an international issue: Vol. Current issues in midwifery series. Quay.

How to suture correctly - Royal College of Midwives. (n.d.).
<http://www.rcm.org.uk/midwives/features/how-to-suture-correctly/>

Kettle, C. and M. D. Raynor. (2010). Perineal management and repair. In Advancing skills in midwifery practice. Churchill Livingstone Elsevier.

Kettle, C., R. K. Hills and K. M. K. Ismail. (2007). Continuous versus interrupted sutures for repair of episiotomy or second degree tears. The Cochrane Database of Systematic Reviews, Issue 4.

Kettle, C...[et al]. (2002). Continuous versus interrupted perineal repair with standard or rapidly absorbed sutures after spontaneous vaginal birth: a randomised controlled trial. The Lancet, 359(9325).

Lee, B. (2002). Are you sitting comfortably? Issues around perineal trauma. RCM Midwives Journal, 5(9).

Lewis, G. (ed), The Confidential Enquiry into Maternal and Child Enquiries (CMACE). (2011). Saving mothers' lives : reviewing maternal deaths to make motherhood safer - 2006-2008 : the eighth report of the confidential enquiries into maternal deaths in the United Kingdom. BJOG an International Journal of Obstetrics and Gynaecology, 118(Suppl 1).

Lewis, Gwyneth, Confidential Enquiry into Maternal Deaths (Great Britain), & Confidential Enquiry into Maternal and Child Health (Great Britain). (2007). Saving mothers' lives: reviewing maternal deaths to make motherhood safer - 2003-2005 : the seventh report of the confidential enquiries into maternal deaths in the United Kingdom. Confidential Enquiry into Maternal and Child Health.

Lieberman, E., & O'Donoghue, C. (2002). Unintended effects of epidural analgesia during labor: a systematic review. American Journal of Obstetrics and Gynecology, 186(5, Supp A), S31-S68. <http://www.sciencedirect.com/science/journal/00029378/186/5/supp/S>

Lockhat, Haseena. (2004). Female genital mutilation: treating the tears. Middlesex University Press.

Lundquist, M...[et al]. (2000). Is it necessary to suture all lacerations after a vaginal delivery? Birth, 27(2).

Macarthur, A.J. and C. Macarthur. (2004). Incidence, severity, and determinants of perineal pain after vaginal delivery: A prospective cohort study. American Journal of Obstetrics and Gynecology, 191(4).

MacArthur, C...[et al]. (2005). Faecal incontinence and mode of first and subsequent delivery: a six-year longitudinal study. British Journal of Obstetrics and Gynaecology, 112(8), 1075-1082. <https://doi.org/10.1111/j.1471-0528.2005.00721.x>

Mackrodt, C...[et al]. (1998). The Ipswich childbirth study: 2. A randomised comparison of

polyglactin 910 with chromic catgut for postpartum perineal repair. *British Journal of Obstetrics and Gynaecology*, 105(4), 441–445.

<http://onlinelibrary.wiley.com/doi/10.1111/j.1471-0528.1998.tb10131.x/pdf>

Mackrodt, C...[et al]. (2001). How cost effective is it to leave perineal skin unsutured. *British Journal of Midwifery*, 9(4), 209–214.

Management of pregnancy, childbirth and the postpartum period in the presence of female genital mutilation: Report of a WHO Technical Consultation. (n.d.).

Management of third and fourth degree perineal tears following vaginal delivery; Guideline no 29. (n.d.).

McCandlish, R...[et al]. (1998). A randomised controlled trial of care of the perineum during second stage of normal labour. *British Journal of Obstetrics and Gynaecology*, 105(12), 1262–1272. <http://onlinelibrary.wiley.com/doi/10.1111/j.1471-0528.1998.tb10004.x/pdf>

McCandlish, R...[et al]. (2003). The HOOP study: robust evidence for perineal care? *British Journal of Midwifery*, 11(3).

Mei-dan, E...[et al]. (2008). Perineal massage during pregnancy: a prospective controlled trial. *Israel Journal of Medical Sciences*, 10(7).

Metcalfe, A...[et al]. (2002). A pragmatic tool for the measurement of perineal tears. *British Journal of Midwifery*, 10(7).

Momoh, Comfort. (2005). *Female genital mutilation*. Radcliffe Pub.

Momoh, Comfort, Barber, Gillian, Basak, Carolyn, & Royal College of Nursing (Great Britain). (2006). *Female genital mutilation: an RCN educational resource for nursing and midwifery staff*. Royal College of Nursing.

Myrfield, K., C. Brook and D. Creedy. (1997). Reducing perineal trauma: implications of flexion and extension of the fetal head during birth. *Midwifery*, 13(4).

National Institute for Health and Clinical Excellence (Great Britain) & National Collaborating Centre for Primary Care (Great Britain). (2006). *Routine postnatal care of women and their babies: Vol. NICE clinical guideline*. National Institute for Health and Clinical Excellence.

National Institute for Health and Clinical Excellence (Great Britain) & National Collaborating Centre for Women's and Children's Health (Great Britain). (2007). *Intrapartum care: care of healthy women and their babies during childbirth: Vol. NICE clinical guideline*. National Institute for Clinical Excellence.

National Institute for Health and Clinical Excellence (NICE). (n.d.).

National Perinatal Epidemiology Unit. (n.d.).

NHS Health Quality. (n.d.).

Nichols, C. M...[et al]. (2006). Anal sphincter defects and bowel symptoms in women with and without recognised anal sphincter trauma. *American Journal of Obstetrics and*

Gynecology, 194(5).

Norton, C...[et al]. (2002). Anal incontinence. In Incontinence (2nd ed.). Plymouth: Health Publication Ltd.

Oberwalder, M., J. Connor and S. D. Wexner. (2003). Meta-analysis to determine the incidence of obstetric anal sphincter damage. British Journal of Surgery, 90(11).

Oboro, V. O...[et al]. (2003). A multicentre evaluation of the two-layered repair of postpartum perineal trauma. Journal of Obstetrics and Gynaecology, 23(1).

Obstetric anal spincter injury: 7 critical questions about care. (n.d.).

Oliver, Paul. (2012). Succeeding with your literature review: a handbook for students: Vol. Open UP study skills. Open University Press.

Power, D., M. Fitzpatrick and C. O'Herlihy. (2006). Obstetric anal sphincter injury: How to avoid, how to repair: A literature review. The Journal of Family Practice, 55(3).

Royal College of Obstetricians and Gynaecologists. (n.d.).

Shorten, A., J. Donsante and B. Shorten. (2002). Birth position, accoucheur, and perineal outcomes: Informing women about choices for vaginal birth. Birth, 29(1).

Sleep, J. and A. Grant. (1987). West Berkshire perineal management trial: three year follow-up. British Medical Journal, 295(6601).

Sleep, J...[et al]. (1984). West Berkshire perineal management trial. British Medical Journal, 289(6445).

Stamp, G., G. Kruzins and C. Crowther. (2001). Perineal massage in labour and prevention of perineal trauma: randomised controlled trial. BMJ : British Medical Journal / British Medical Association, 322(7297).

Steen, M., & Marchant, P. (2001). Alleviating perineal trauma the APT study. RCM Midwives Journal, 4(8), 256-259.

Steen, M...[et al]. (2000). A randomised controlled trial to compare the effectiveness of ice-packs and epifoam with cooling maternity gel pads at alleviating postnatal perineal trauma. Midwifery, 16(1).

Sultan, A. H. and K. Kettle. (2009). Diagnosis of perineal trauma. In Perineal and anal sphincter trauma: diagnosis and clinical management. Springer.

Sultan, Abdul H., Thakar, Rane, & Fenner, Dee E. (2009). Perineal and anal sphincter trauma: diagnosis and clinical management. Springer.

Symon, A. (2008). Third degree tears: the three stage negligence test. British Journal of Midwifery, 16(3).

Thacker, S. B. (2000). Midline versus mediolateral episiotomy. BMJ : British Medical Journal / British Medical Association, 320(7250).

The Birth Trauma Association. (n.d.).

The Royal College of Midwives. (n.d.).

Wickham, S. (2007). Research unwrapped: Preventing perineal damage in childbirth. *The Practising Midwife: The Voice of Clinical Midwifery*, 10(2), 37–39.

Williams, A...[et al]. (2005). Women's experiences after a third-degree obstetric anal sphincter tear: A qualitative study. *Birth*, 32(2).

Williams, A...[et al]. (2006). How to repair an anal sphincter injury after vaginal delivery: results of a randomised controlled trial. *British Journal of Obstetrics and Gynaecology*, 113(2).

Wilson, A. E. (2012). Effectiveness of an educational programme in perineal repair for midwives. *Midwifery*, 28(2), 236–246. <https://doi.org/10.1016/j.midw.2011.02.011>

World Health Organisation. (n.d.).

World Health Organisation Study Group on Female Genital Mutilation and Obstetric Outcome. (2006). Female genital mutilation and obstetric outcome: WHO collaborative prospective study in six African countries. *The Lancet*, 367(9525).